

CANNA TERRA

GROW SCHEDULE



	VEGETATIVE PHASE				GENERATIVE PHASE						
	ROOTING	I	II		I			II	III	IV	
DAYS AND WEEKS	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
LIGHT HOURS	18	18	12	12	12	12	12	12	12	12	12
	ml/gal										
CANNA TERRA VEGA	15.5	20.3	23.6	23.6							
CANNA TERRA FLORES					25.1	25.1	25.1	25.1	25.1	16.5	
CANNA RHIZOTONIC	15.0	7.5	7.5	7.5	2.0	2.0	2.0	2.0	2.0		
CANNAZYM		9.5	9.5	9.5	9.5	9.5	9.5	9.5	9.5	9.5 - 18.9*	9.5 - 18.9
CANNABOOST**			7.6	7.6	7.5 - 15.0	7.5 - 15.0	7.5 - 15.0	7.5 - 15.0	7.6 - 15.1	7.5 - 15.0	7.5 - 15.0
CANNA PK 13/14								5.7			
CANNA CALMAG AGENT***	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0	0 - 4.0
	mS/cm										
EC WATER***	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
EC NUTRITION	1.0	1.3	1.5	1.5	1.5	1.5	1.5	1.9	1.5	1.0	
EC TOTAL	1.4	1.7	1.9	1.9	1.9	1.9	1.9	2.3	1.9	1.4	0.4

Rooting:
Start/rooting;
hydrate substrate.

Vegetative phase I:
Plants develop in volume.

Vegetative phase II:
Up to growth stagnation after fructification or appearance of the formation of flowers.

Generative phase I:
Flowers or fruits develop in length. Growth in height achieved.

Generative phase II:
Development of the volume (breadth) of flowers or fruit.

Generative phase III:
Development of the mass (weight) of flowers or fruit.

Generative phase IV:
Flowers or fruit ripening process.

This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase depends on crop species, cultivar, desired plant size and/or plants per m².

- * Double CANNAZYM dosage to 19 ml/gallon, if substrate is reused.
- ** Increase to a maximum of 15 ml/gallon through flowering for extra strength.
- *** Adjust CANNA CALMAG AGENT application rate to achieve an ECwater of 0.2-0.4 mS/cm. This example schedule assumes that the ECwater has been adjusted with Calmag or naturally contains an EC of 0.4 mS/cm.

To set your plants up very early on as a cutting or seedling, check out CANNA Start. This is to be used prior to commencing the grow guide above. For a longer flowering phase, repeat the recommended dosages from Generative Phase I (W3-W5). Add CANNA PK 13/14 in the period 3-4 weeks before harvest.

Recommended pH is between 5.5 and 6.2.

